

Alexander E. Weber, MD
 Sports Medicine Surgery
 Team Physician USC Athletics and LA Kings
Appointments: 855-SC-SPORT
Office: 818-658-5921
Cellular: 518-928-8389
Email: weberae@usc.edu
Website: www.AlexWeberMD.com



MENISCAL ALLOGRAFT TRANSPLANTATION REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-2 weeks	Heel touch weight bearing with crutches	Locked in full extension for sleeping and all activity Off for exercises and hygiene	0-90° when non-weight bearing	Heel slides, quad sets, patellar mobs, SLR, SAQ** No weight bearing with flexion >90°
PHASE II 2-8 weeks	2-6 weeks: Heel touch WB with crutches 6-8 weeks: Progress to full WB	2-8 weeks: Locked 0-90° Off at night Discontinue brace at 8 weeks	As tolerated	Addition of heel raises, total gym (closed chain), terminal knee extensions** Activities w/ brace until 6 weeks; then w/o brace as tolerated No weight bearing with flexion >90°
PHASE III 8-12 weeks	Full	None	Full	Progress closed chain activities Begin hamstring work, lunges/leg press 0-90°, proprioception exercises, balance/core/hip/glutes Begin stationary bike
PHASE IV 12-20 weeks	Full	None	Full	Progress Phase III exercises and functional activities: single leg balance, core, glutes, eccentric hamstrings, elliptical, and bike Swimming okay at 16 wks
PHASE V > 20 weeks	Full	None	Full	Advance to sport-specific drills and running/jumping once cleared by MD

*Brace may be removed for sleeping after first post-operative visit (day 7-10)

**Avoid any tibial rotation for 8 weeks to protect meniscus