Alexander E. Weber, MD

Sports Medicine Surgery

Team Physician USC Athletics and LA Kings

Appointments: 855-SC-SPORT

Office: 818-658-5921 Cellular: 518-928-8389 Email: weberae@usc.edu

Website: www.AlexWeberMD.com



## MPFL ALLOGRAFT RECONSTRUCTION REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-2 weeks	As tolerated with brace	On at all times during day and while sleeping** Off for hygiene	0-90°	Heel slides, quad and hamstring sets, patellar mobs, gastroc stretch, SLR in brace, modalities
PHASE II 2- 6 weeks	As tolerated with brace	Discontinue at 6 weeks if able to obtain full extension w/o lag*	Maintain full extension and progress flexion	Progress weight bearing flexibility, begin toe raises and closed chain quad work Begin floor-based core
				and glutes work, balance exercises, hamstring curls and stationary bike
PHASE III 6 weeks - 4 months	Full	None	Full	Advance closed chain quads, progress balance, core/pelvic and stability work
				Begin elliptical, in-line jogging at 12 weeks under PT supervision
PHASE IV 4-6 months	Full	None	Full	Progress flexibility/strengthening, progression of function: forward/backward running, cutting, grapevine, initiate plyometric program and sport- specific drills @ 16 wks
				Return to play as tolerated after 16 weeks post-op when cleared by MD

<sup>\*</sup>Brace may be changed to patellar-stabilizing brace (Hely-Weber, Shields, or equivalent) after first post-operative visit

<sup>\*\*</sup>Brace may be removed for sleeping after first post-operative visit (day 7-10)